

## 钾离子通道多聚体结构域蛋白 20 抗体

产品货号： mlR16931

英文名称： KCTD20

中文名称： 钾离子通道多聚体结构域蛋白 20 抗体

别名： BTB/POZ domain-containing protein KCTD20; C6orf69; dJ108K11.3; KCD20\_HUMAN; KCTD20;  
Potassium channel tetramerisation domain containing 20.

研究领域： 细胞生物 神经生物学 信号转导 通道蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 47kDa

细胞定位 : 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KCTD2:181-280/419

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.



**PubMed :** PubMed